

# Epigentic Changes In Endometrial Cancer -: An Important Area Of Research

# Zubiya Ahsan<sup>1</sup>,Neeraj Vaishnav<sup>2</sup>,Debasray Saha<sup>1</sup>Runjhun Mathur<sup>3</sup>, Abhimanyu Kum ar Jha<sup>1</sup>\*

1. Department of Biotechnology, Faculty of Life Sciences, Institute of Applied Medicines and Research, Ghaziabad, Uttar Pradesh, India.

2. Department of Biotechnology, University of Rajasthan, Jaipur, Rajasthan, India.

3.Dr. A.P.J Abdul Kalam Technical University, Lucknow, India.

\*- Corresponding Author. Email address- abhimanyujha630@gmail.com

#### ABSTRAC T

Endometrial cancer is the most commonly diagnosed gynaecological cancer. Based on differences in clinic pathologic characteristics, there are two subtypes of endometrial carcinoma Type 1 and Type 2. Multiple risk factors(Obesity, Use of an intrauterine device, Age, Diet and Exercise, Type 2 Diabetes) are developing endometrial cancer. Likewise, epigenetics is the study of alteration gene and modulated gene expression with alteration in DNA sequences. In mammals, epigenetic modifications include DNA methylation, histone modifications, and miRNA.

Epigenetics alterations are associated with promoter methylation of DNA such as hMLH1( MutL homolog1), hM SH2(M uts homolog2), hM SH3, hM SH6 and hPM S2 and demethylation of tumor suppressor genes such as PTEN, RASSF1A, APC, HOPX, HOXA10, and MGMT. Promoter methylated genes(Phosphatase and tensin homolog Ras association domain family 1 isoform, Antigen-presenting cell, Homeobox Only Protein Homeobox, Homeobox protein Hox-A10, O6 methylguanine DNA methyltransferase)are related to cell proliferation, development, and differentiation. This study concludes the effect of promoter methylation genes on genetic and epigenetic modifications and reveals how these genes can act as a biomarker and can improve health outcomes.



#### **INTRO DUCTION**

Endometrial cancer is the most commonly diagnosed gynaecological cancer. It is the 4<sup>th</sup> most common cancer in women[1]. Endometrial means that the cancer starts in the lining of the womb. This lining is called the endometrium. About 95% of endometrial cancers are adenocarcinomas. Adeno means that the cells that have become cancerous. They are the cells of glandular tissue. So for the most common type of womb cancer, the cancer is in the glands of the endometrium. Based on differences in clinic pathologic characteristics, there are two subtypes of endometrial carcinoma. Type 1 cancers are the most common type. They are usually endometrioid adeno carcinomas. Type 2 cancers include uterine serous carcinomas and clear cell carcinomas. Type1 cancers are linked to excess estrogen whereas Type 2 cancers are not linked to excess estrogen. Type I tumors are usually well-differentiated, and most patients present with early-stage of the disease and have a favourable prognosis. In contrast, type II, non-endometrioid endometrial carcinoma, is more common in older postmenopausal women, often poorly differentiated Type I tumors commonly have neardiploid karyotypes, microsatellite instability and mutations in PTEN, K-RAS and CTNNB1 ( $\beta$ -catenin) genes; whereas type II tumors are characterized more often by p53 mutation, overexpression of Her-2/neu and an aneuploid karyotype[2]. Although multiple risk factors affect the risk of developing endometrial cancer, such as obesity, the use of an intrauterine device (IVD), age, hormone factors, estrogen therapy these factors have been identified the understanding of the etiologies of the two subtypes of endometrial cancer continues to evolve. Endometrial cancer is a complex disease driven by epigenetic alterations, abnormal genetic as well as environmental factors, As a common molecular alteration in human neoplasia, epigenetics is the study of heritable phenotype changes that do not involve alterations in the DNA sequence [3]. Epigenetic changes are equally responsible for genetic changes in the development and progression of cancer. Epimutation change can assist determine whether factors are turned on or off controlling the production of macromolecules in particular cells [4].

These epigenetic processes are necessary which regulate the normal functions of the cell during all the stages, including development, differentiation and facilitate adaptation to environmental changes, such as nutritional variation of exposure to cigarette smoke, chemicals, radiation, and hormones[5].



#### MECHANISM OF EPIGENETIC MODIFICATIONS

Epigenetic modifications can be defined as heritable alterations in gene expression and cellular function without changes to the original DNA sequence. The processes responsible for epigenetic regulation are DNA methylation, modifications in chromatin, and noncoding RNA(miRNA).

#### DNA Methylation

The DNA of most organisms is modified by a post-replicative process which results in three types of methylated bases in DNA: C5-methylcytosine, N4-methylcytosine, N6-methyladenine, This modification is called DNA methylation. It is an epigenetic mechanism used by cells to control gene expression. Some mechanisms exist to control gene expression in eukaryotes, but DNA methylation is a commonly used epigenetic signaling tool that can fix genes in the "off" position. Addition of methyl group to the C-5 position of cytosine residues. Most cytosine methylation occurs in the sequence context 5'CG3'. This occurs almost exclusively at cytosines that are followed immediately by a Guanine- CpG Dinucleotide. Methyl groups are transferred from S-adenosyl methionine in a reaction catalyzed by DNA methyltransferases (DNMT) or methylases. SAM is then converted to SAH (S-adenosyl homocysteine).

#### **Histone Modification**

Histones are subject to a wide variety of posttranslational modifications including but not limited to, lysine acetylation, lysine and arginine methylation, serine and threonine phosphorylation and lysine ubiquitination and sumoylation. These modifications occur primarily within the histone amino-terminal tails protruding from the surface of the nucleosome as well as on the globular core region.

Histone modifications are proposed to affect chromosome function through at least two distinct mechanisms. The First mechanism suggests modifications may alter the electrostatic charge of the histone resulting in a structural change in histones or their binding to DNA. The Second Mechanism proposes that these modifications are binding sites for protein recognition modules, such as the bromodomains or chromodomains, that recognize acetylated lysines or methylated lysine, respectively.



#### Mi RNAs

miRNAs or ncRNA represent small RNA molecules encoded in the genomes of plants and animals. These highly conserved 22 nucleotides long RNA sequences regulate the expression of genes by binding to the 3'-untranslated regions (3'-UTR) of specific mRNAs. A growing body of evidence shows that miRNAs are one of the key players in cell differentiation and growth.

## PROMOTER HYPERMETHYLATION OF GENES IN ENDOMETRIAL CANCER

#### DNA mismatch repair gene in endom etrial cancer.

Hereditary non-polyposis colorectal cancer (HNPCC), or Lynch Syndrome, is one of the most common familial cancer syndromes. The molecular basis for HNPCC is a defect in a DNA mismatch repair gene, usually a germline mutation **in hMLH1, hMSH2, hMSH3, hMSH6, and hPMS2.** A defective DNA mismatch repair causes microsatellite in stability and, a high risk of the colon as well as endometrial cancers. The hMLH1 and hMSH2 mutations are rare (less than 10%) in sporadic endometrial cancers in comparison to the MSI+ phenotype. However, reduced protein expression of hMLH1 and other mismatch repair genes is a common finding in endometrial cancers. The study of these MMR proteins has revealed that recognition of the base-base mismatches and deletion/insertion loops is performed by a heterodimer of either MSH6 and MSH2 or MSH2 and MSH3. The MSH2 and MSH3 are heterodimers preferentially recognizes insertion and deletion loops. Consequently, cancers arising with a loss of MSH6 function display microsatellite instability.

A heterodimer of MLH1-PMS2 operates as a molecular matchmaker and is involved in executing the repair of mismatches. Promoter methylation is associated with a lack of expression of hMLH1 in human cancers and mismatch repair-defective human tumor cell lines.

#### Methylation of steroid receptor genes in endometrial cancer.

Endometrium is highly responsive to hormonal stimuli. The cyclic production of estrogen and progesterone during the menstrual cycle, and declining sex steroid hormone levels after menopause are directly correlated with endometrial proliferation and morphological changes.



There are several risk factors that directly or indirectly affected by estrogen. Several studies have been done, evaluated the association of promoter methylation of the estrogen receptor (ER) and progesterone receptor (PR) genes in endometrial cancer.

**Estrogen receptors** (ERs) are group 0f proteins found inside the cells. Genes are located at chromosome 6q25.1, have a CpG island in its promoter and exon 1 regions. Commonly they are receptors but activated by the hormone estrogen. Two classes of ER are there, One is nuclear receptor family of intracellular and membrane estrogen and the other is, once activated by estrogen, the ER can translocate into the nucleus and perform the function of DNA binding. This is the proliferative phase of the endometrium. It strongly binds to SHBG and albumin.

**Progesterone receptors** (PR) is a secondary phase of the endometrium. It binds to corticosteroid-binding globulin and albumin. It is a nuclear receptor and protein found inside cells belong to subfamily 3, group C and member 3. It is encoded by a single PGR gene residing on chromosome 11 q22 in humans. They are ligand-activated transcription factor members of the steroid hormone.

#### Methylation of tumor suppressor genes in endometrial cancer.

Promoter hypermethylation of tumor suppressor genes is a major event in the origin of many cancers. Some studies have been done and a list of tumor suppressor genes frequently hypermethylated in endometrial cancer are PTEN, RASSF1A, APC, HOPX, HOXA10, and MGMT.

#### PTEN

The phosphatase and tensin homolog (PTEN, also known as MMAC1/TEP1) gene on chromosome 10q23.3[5]. It provides instructions for making an enzyme. It is characterized as a tumor suppressor, which means that it regulates cell division by keeping cells from growing and dividing too rapidly or in an uncontrolled way. Promoter hypermethylation, as the alternative mechanism of PTEN allelic inactivation and occurred in sporadic colorectal tumors with microsatellite instability[6].



# RASSF1A

The Ras association domain family 1 isoform(RASSF1A) neoplastic genes were regulated like cell cycle suppression, programmed cell death and here dity instability that shows a vital role in suppressing Ras-mediated oncogenesis[7]. Moderation of each allele of a neoplasm inhibitor gene is necessary for carcinoma. RASSF1A is the most commonly hypermethylated neoplasm suppressor genes in human tumors and may assist as a biomarker for disease detection[8]. Natural compounds such as curcumin and resveratrol inhibit the activity of DNA methyltransferases.

#### APC

Adenomatous polyposis coli (APC), a tumor suppressor gene, regulates  $\beta$ -catenin in the Wntsignaling path way[9]. It provides instructions for making a protein that keeps cells from growing and dividing rapidly in an uncontrolled manner. This helps to ensure the number of chromosomes in the cell is correct following cell division. APC gene promoter methylation has been observed around 20%-45% in endometrial cancer[10].

#### MGMT

The 06 methylguanine DNA methyltransferase (MGMT) is a polymer repair macromolecule that protects the human regulating from agents and tumor conduct of endogenous cancer[11]. Also, it induces protection to alkylating agents and removes alkyl adducts from the O6 position of guanine in DNA.

#### **P16**

The p16 gene additional called CDKN2A encodes p16INK4A that suppresses the CDK45: Cyclinand CDK6: cyclin D complex es[12]. Hypermethylation of the p16 neoplasm suppressor gene and has effects on transcriptional down-regulation or silencing, this has been shown as one of the vital structures of p16INK4a gene inactivation in varied kinds of carcinoma such as colon cancer, thyroid cancer, and cervical cancer[13].



### RAR<sub>\$2</sub>

Retinoic Acid Receptor (RAR $\beta$ 2) is known as a cancer inhibitor gene by correlating with retinoic acid Expression of retinoic acid receptor  $\beta$  (RAR $\beta$ ) is defined to be absent or disease controlled in cancer [14]. Epigenetic silencing by promoter hypermethylation ends up in retinoic acid therapy failure in breast carcinoma [15].

#### **CONCLUSION**

Recent Advancements in the field of epigenetics, help in understanding the role of epigenetic alterations in normal cellular processes and abnormal changes leading to endometrial carcino genesis. It is been observed that DNA methylation plays a key role in understanding the promoter methylation genes in endometrial cancer. By understanding the promoter methylation genes some of epigenetic changes can be studied in endometrial cancer. promoter methylation thesegenes can act as a biomarker. This canact as an important tool against endometrial cancer in the future all over the world

#### **REFERENC ES**

1. American Cancer Society. (2009) Cancer Facts and Figures 2009. www.cancer.org.

2. Prat J, Gallardo A, Cuatrecasas M, Catasús L. Endometrial carcinoma: pathology and genetics. Pathology 2007; 39:72-87.

3. Ledford H (October 2008). "Language: Disrupted definitions". Nature 455(7216):1023-8.

4.Illin gworth R, kerr A Desousa D, Jorgensen H, Ellis P, Stalker J, et al. A novel CpG island set identifies tissue-specific mmethylation at development gene loci.PLoS Biol.2008;6:e22.

5. Nelen MR, Padberg GW, Peeters EA, Lin AY, van den Helm B, Frants RR, et al. Localization of the gene for Cowden disease to chromosome 10q22-23. Nat Genet 1996; 13:114-6.



6. Goel A, Arnold CN, Niedzwiecki D, Carethers JM, Dowell JM, Wasserman L, et al. Frequent inactivation of PTEN by promoter hypermethylation in microsatellite instabilityhigh sporadic colorectal cancers. Cancer Res 2004; 64:3014-21.

7. Donninger H, Clark J, Rinaldo F, Nelson N, Barnoud T, Schmidt ML, et al. The RASSF1A tumor suppressor regulates XPA-mediated DNA repair.Mol Cell Biol2015;35(1):277-87

8. Garzon R, Calin GA, et al, MicroRNAs in cancer. Annu Rev Med 2009;60:167-179(PubMed 19630570)

9. Fodde R. The APC gene in colorectal cancer. Eur J Cancer 2002; 38:867-71.

10. Banno K, Yanokura M, Susumu N, Kawaguchi M, Hirao N, Hirasawa A, et al. Relationship of the aberrant DNA hypermethylation of cancer-related genes with carcinogenesis of endometrial cancer. Oncol Rep 2006; 16:1189-96.

11. Fang M, Chen D, Yang CS. Dietary polyphenols may effect DNA methylation. J Nutr, 2007;137(Suppl1):223S-228S.

12. Duthie SJ. Epigenetic modifications and human pathologies: Cancer and CVD. Proc Nutr Soc. 2011;70(1):47-56

13. Hayashi K, Yokozaki H, Naka K, Yasui W, Lotan R, Tahara E. Overexpression of retinoic acid receptor beta induces growth arrest and apoptosis in oral cancer cell lines. Jpn J Cancer Res. 2001;92:42-50

14. Jha AK, Nikbakht M, Parashar G, et al. Reversal of hypermethylation and reactivation of the RARbeta2 gene by natural compounds in cervical cancer cell lines. Folia Bio(Praha). 2010;56:195-200

15. Mittag F, Kuester D, Vieth M, Peters B, DAPK promoter methyaltion is an early event in colorectal carcinogenesis. Cancer Lett. 2006;240:69-75