

Maternal and Newborn Mortality: Community Opinions on Why Pregnant Women and Newborns Are Dying In Natikiri, Mozambique.

Belo C,¹ Pires P,² Josaphat J,³ Siemens R,⁴ Rooke E,⁵ Spence-Gress C.⁶

ABSTRACT

Background and objective: maternal and neonatal mortality rates in Mozambique are high, due to insufficient numbers of qualified health workers, lack of equipment and materials, referral system deficiency, difficulties to access health services and gender issues. This study assesses the barriers to health care access, regular attendance at ante natal consultations, institutional delivery and postnatal and neonatal follow-up. This is part of the baseline study for an implementation research project to reduce maternal and neonatal mortality in Natikiri, Nampula, Mozambique.

Methods: descriptive mixed study with two components: 1) data analysis from primary sources (interviews and focus group discussions with community members, health professionals) in the catchment area of Marrere health center and hospital, in Natikiri; 2) data analysis of secondary sources (national and international literature).

Results: 300 people were surveyed and 11 focal group discussions were held. Respondents were asked why they thought pregnant women and newborns were dying in their community. Local community members and health professionals some reasons:

1. Long walking distances required to reach health services.
2. Unsafe travel conditions for women.

¹ Project leader, conception, data collection and interpretation, final approval of the version to be published; MD Master in Occupational Health, Health Sciences Faculty Dean, Lúrio University, Nampula, Mozambique.

² Study protocol conception and design, data collection, analysis and interpretation, article draft, final approval of the version to be published; Family and Community Medicine Specialist, Lecturer, Health Sciences Faculty, Lúrio University, Nampula, Mozambique.

³ Study protocol conception and design, data collection, treatment, analysis and interpretation, article draft, final approval of the version to be published; Nurse, Health Services Management Specialist, Master in Epidemiology, Lecturer, Health Sciences Faculty, Lúrio University, Nampula, Mozambique.

⁴ Study protocol conception, data collection and interpretation, translation to English, final approval of the version to be published; MD Paediatrician, Lecturer, University Saskatchewan, Saskatoon, Canada.

⁵ Study protocol conception, data collection and interpretation, final approval of the version to be published; MD Family Medicine Specialist, Lecturer, University Saskatchewan, Saskatoon, Canada.

⁶ Study protocol conception, data treatment, analysis and interpretation, final approval of the version to be published; **MD Family Medicine Specialist**, Lecturer, University Saskatchewan, Saskatoon, Canada.

3. Poor treatment in health facilities including illicit payments and bribes.
4. Long waits to be attended to.
5. Poor training and lack of knowledge of health professionals.
6. Health professionals neglecting patients and not giving family centered care.
7. Teenage pregnancies and short spacing of pregnancies.
8. Inability of women to make informed decisions about family planning.
9. Home births without trained support and traditional methods of treatment.
10. Myths and cultural taboos about pregnancy and newborn care.
11. Women and community limited knowledge about women and adolescent girls' health.
12. Need of "mother's house waiting" near the hospital for pregnant women.
13. Weak government policies and little funding to support maternal and child health care.

Discussion: literature review identified several factors causing delay in pregnant women and newborns' appropriate care. These poor quality determinants on primary and secondary health care, for pregnant and newborn, can be grouped

into three delays: (1) the decision to seek care by pregnant woman and woman who have delivered; 2) accessing and arriving at the health center; 3) receiving quality health care.

Conclusion: local community members and health professionals were asked to state what they thought would be the best way to intervene. These ideas were then discussed further at a conference with health professionals and government representatives. Six intervention strategies to address identified problems were decided on and will be the basis for the ongoing implementation research project. They were:

1. Expanding family planning especially with adolescents.
2. Community based transport system for pregnant women.
3. Strengthening maternal and child health services by training maternity personnel in obstetrical emergency care and neonatal resuscitation.
4. Providing four quality prenatal visits.
5. Providing quality cesarean deliveries.
6. Supporting the Mozambican Government's campaign against bribery.

Keywords: access, prenatal, consultation, pregnancy, puerperium, newborn, Mozambique.

1. Introduction

This study carried out by Lúrio University (UniLúrio) Health Sciences Faculty (HSF) in partnership with Nampula Provincial Health Directorate (NPHD), Marrere Hospital (MH) and the University of Saskatchewan, Saskatoon, Canada constitutes part of the baseline evaluation for an implementation research on maternal and newborn health.

Every day, about 800 women die from preventable causes related to pregnancy and childbirth. Almost three million newborn babies die each year and 2.6 million babies are stillborn.¹

The World Health Organization (WHO) defines maternal mortality as the death of a woman during pregnancy or up to 42 days after giving birth, irrespective of the duration and place of the pregnancy, due to any cause related or aggravated by pregnancy or its management. Currently maternal and neonatal mortality rates in Mozambique are unacceptably high: 4,800 maternal deaths during the year 2013.²

The complications responsible for almost 75 % of maternal deaths in the

world are: severe bleeding and infections (usually after childbirth), high blood pressure during pregnancy (preeclampsia and eclampsia), other childbirth and unsafe abortion complications.³

In Mozambique, about 43 % of maternal deaths occur during childbirth and up to 24 hours later, 76 % of these deaths were due to direct causes and 24 % to indirect causes. Among main causes of death are uterine rupture (17 %), postpartum haemorrhage (14 %), preeclampsia and eclampsia (13 %), Acquired Human Immunodeficiency Syndrome (AIDS) (12 %) and puerperal sepsis (11 %). AIDS appears as the first indirect cause of maternal death and the fourth leading cause. In primary health care (PHC) units death occurs more frequently before the woman reaches the first two hours of hospitalization, showing the precarious conditions and women's late arrival to the health center (HC).⁴

The main causes of neonatal morbidity and mortality are prematurity and low birth weight, asphyxia, sepsis, pneumonia, human immunodeficiency virus (HIV), malaria, diarrhea, syphilis and other congenital infections. In addition to these factors, the low frequency of institutional delivery, (54 % in 2011), the reduced quality and

quantity of antenatal consultations (ANC) also contribute to morbidity and mortality.^{5,6}

In Mozambique, among other determinants of maternal and neonatal deaths, are the shortage of qualified personnel in the HC, poor quality and quantity of materials and equipment, low quality care, deficiency in referral system, long travel distances to the HC, lack of transportation, poor communication between health professionals (HP) and the community, and gender issues such as the low decision-making power of women and low literacy levels.^{7,8}

These factors can be grouped using the three-delay model: 1) delay in the decision to seek appropriate maternal or neonatal health care; 2) delay in arrival to the HC; 3) delay in receiving timely and appropriate obstetric or neonatal emergency care.⁹

The national health system (NHS) covers 40 % of the population with hospital care and 60 % in PHC. The remaining population is covered by a community network composed of traditional midwives (TMW) and traditional health practitioners (THP).

A study in Mali showed that organized participation of TMW improved the

access of women to ANC, institutional delivery, neonatal follow-up and children vaccination.¹⁰

Well equipped maternities with trained personnel are key to provide skilled birth attendants and deal with obstetric complications. However there is a persistence of inequities in HC distribution in the country: about half of pediatricians and obstetricians are concentrated in Maputo, the Mozambican capital.

The proportion of births in Mozambique in HC with trained birth attendants increased from 48 % in 2003 to 55 % in 2011 but neonatal mortality has declined more slowly than infant and child mortality. This puts the country still far from achieving the annual decline needed to achieve the Sustainable Millennium Goals (SMG).¹¹

Considering the importance of delivery and birth care for maternal and neonatal health, it is necessary to invest in actions with a positive impact on this reality. The best potential to modify the situation is in the puerperium and immediate postpartum (the first 24 hours after childbirth) where 24 to 45% of neonatal and 45% of maternal deaths occur.^{12, 13}

Several programs have been developed in Mozambique to reduce maternal and

neonatal mortality. One is the distribution to pregnant women of the Women's Health Handbook.¹⁴ The Community Health Program, implemented through Local Health Councils, aimed to reduce family and community barriers to access ANC and increase social mobilization. Another program focused on sexual and reproductive health (SRH) barriers, by building "Mother's Waiting Homes" to house mothers near the HC while awaiting delivery, reviewing the abortion law, using information from both men and women and extending the coverage of SRH services.¹⁵

The creation of Hospital Co-management Committees, participated by HP and managers and community members who work together in the planning, implementation, follow-up and evaluation of activities, including analysis and decision-making in the HC and the community, aims to improve health services' performance.¹⁶

The Model Maternity Program aims to improve the quality of delivery and the humanization of health services to women and children, but to date with a reduced implementation.¹⁷

The objective of this study is to evaluate the barriers to access and adherence to

ANC, institutional birth and follow-up during puerperal and neonatal periods in the HC and MH. The secondary objectives were: (a) analyze local community' perception about pregnancy, childbirth, care during puerperal and neonatal periods; b) assess HP providing Maternal and Child Health (MCH) assistance' knowledge in HC and MH, regarding barriers to access and attendance; c) evaluate TMW and THP knowledge regarding pregnancy, childbirth and care during the puerperal and neonatal periods.

2. Methods

Descriptive mixed-methods research, using quantitative and qualitative data collection from two sources:

1) data analysis from secondary sources (scientific communications, HC and MH reports and programs, scientific articles, national policy and strategy reports, population and health surveys, population census, national and international data).

2) data analysis from primary sources (interviews with provincial department members responsible for MCH, district medical director, MH and HC workers in the area of MCH, TMW, THP, Natikiri community members including women of childbearing age, pregnant women,

postpartum women, adolescents, fathers, elderly women and community leaders).

Interviews used semi structured questionnaires, pre-tested and approved, addressing questions about pregnancy, childbirth and care during puerperium and neonatal period.

Focal group discussions (FGD) were conducted with MCH workers, TMW, THP, community members including women of childbearing age, pregnant women, post-partum women, adolescents, parents, grandparents and community leaders residing in the same area with two questions: why are women and children dying during pregnancy and the newborn period in your community? Secondly, what are possible solutions to the problems identified?

Statistics and reports regarding numbers of deliveries and complications occurring at the HC and MH (hospital for Natikiri district) were collected.

The study was approved by the Institutional Committee on Health Bioethics of Lúrio University and the Behavioral Ethics Board at the University of Saskatchewan.

A representative sample of women of childbearing age was calculated using the Epi Info™ 7.2 program considering the size of the target population, the

expected frequency, with a margin of error of 10% and a 95% confidence interval. The sample size for heads of households was the same as for women of childbearing age: heads of households were considered as partners of women of childbearing age. The same sample size was considered for older women (above 45 years): considered as mothers / mothers-in-law of women of childbearing age. A total of 125 THP operate in the intervention area and the choice of a representative sample used Epi Info™ 7.2 program with the expected frequency of 50%, with a 10% margin of error and a 95% confidence interval. TMW, HP and community leaders willing to participate in the interviews were included.

The estimated sample size for the interviews was 328 people (72 women of childbearing age, 72 male heads of households, 72 elderly women, 54 THP, 32 TMW, 14 HP and 12 community leaders.

11 focus groups were formed with at least 5 participants and a maximum of 12 participants in each group: two groups of women of childbearing age over the age of 18, one for mothers under the age of 18 (adolescents), one for heads of households, one for elderly

women over 45, one for community leaders, one for THP, two of TMW and two of HP.

Participants included had the following criteria:

1. Women of reproductive age, pregnant women, puerperal women over 17 years of age.
2. Adolescents of reproductive age, pregnant, puerperal women aged 17 years or less.
3. Head of household (parents, mothers-in-law and grandparents).
4. Community leaders.
5. HP.
6. THP, TMW.
7. Those who are able and willing to give informed consent.

Different variables and information supports were used for each target group in interview sheets with multiple choice questions and answers (adapted Likert scale):

- 1) Woman of childbearing age: age, residence, household number, level of schooling, number of pregnancies, number of institutional births, number of abortions; trimester of pregnancy in which the first ANC was performed; number of ANC visits during the last pregnancy; number of follow-up visits for the child during the first year; reason

for delay in prenatal and non-institutional delivery (42 questions).

- 2) Elderly woman (mother in law or grandmother): age, residence, number in household, level of education; cause invoked for delay in prenatal and non-institutional delivery (41 questions).

- 3) Head of household: age, residence, number of people in household, level of education, occupation, number of children (41 questions).

- 4) HP: age, gender, profession, position, reasons for delayed ANC visits and non-institutional delivery (40 questions).

- 5) THP / TMW: age, gender, residence, reasons for delayed ANC visits and non-institutional delivery (41 questions).

- 6) Community leader: age, gender, area of residence, level of schooling, reason for delayed ANC visits and non-institutional delivery (41 questions).

- 7) Hospital indicators and statistics: maternal and newborn statistical data collection sheets.

Marrere, in the Administrative Post of Natikiri, City and District of Nampula, is located to the west of the city in plateau zone, mainly of sedimentary soil with granite outcrops and is crossed by six Rivers (Namialo, Marrere, Muepelume, Mussarne, Monapo, Mutivazi). Climate is tropical humid (rainfall > 1,000 mm /

year). Urbanization is deficient (minimal roadways, water supply, domestic sanitation and solid waste collection system) and the population is mostly dispersed, residing in precarious housing of traditional model.

It is estimated that childbearing age women population reaches 10,088 inhabitants (18 % of the total population).

Traditional authorities play an important role in social organization and culture imposes rules and taboos related to pregnancy, childbirth and newborns, often with a negative impact on their health.

MH is located in Marrere subdivision of Natikiri, 12 km from the central part of the city. It is associated with an HC, near a secondary school and refers more difficult health issues to Nampula Central Hospital (NCH) in Nampula city. The radius of the catchment area of MH is about 15 km, with an estimated population of 56,025. It provides pediatric, maternity, emergency, general medicine, basic surgery, radiology, pharmacy, blood bank, vaccination program, and administrative and support services. It has a total of 140 in patient beds. It serves as a site for nursing and medical students training periods and,

participates in the training of allied HP (students of the Health Sciences Institute of Nampula). It is the referral hospital for patients with tuberculosis in the northern part of the country. In 2015, 1,560 births were attended. There are 12 beds for obstetrics and gynecology services. The HP for the care of women and newborns consist of four general practitioners, eight nurses with maternity training, six with basic maternity training and four elementary midwives. There are no Obstetrician / Gynecologists at the hospital.

The interviews and focus groups were done using a Macua (local language) translator.

Data collection focused on the following areas:

A) Challenges faced by pregnant women in the community, preventing them from accessing and joining ANC visits and monitoring during the puerperium and neonatal period.

B) Challenges faced by pregnant women on the way to the HC / MH, preventing them from accessing and joining ANC's and follow-up during the puerperium and neonatal period.

C) Challenges faced by pregnant women in the HC / MH, preventing them from accessing and joining ANC visits and

follow-up during the puerperium and neonatal period.

D) What is there in communities, HC and MH that is working well and what should be improved in relation to pregnancy, childbirth and neonatal period?

E) Existing traditional knowledge in the community related to pregnancy, childbirth and birth.

F) Perception of mothers, pregnant women, postpartum women on ANC and follow-up during the puerperium and neonatal period.

G) Perception of parents and community leaders about ANC visits and follow-up during the puerperium and neonatal period.

H) Perception of TMW and THP on ANC visits and follow-up during the puerperium and neonatal period.

I) HP perception about ANC visits and follow-up during the puerperium and neonatal period.

Data were collected by a team of research assistants composed of students of the UniLúrio medical and nursing courses who speak the local language (Macua) after theoretical and practical

training, including pre-testing of data collection instruments.

Participants were interviewed in their most comfortable language (Portuguese or Macua) to improve understanding of the questions and the elaboration of the answers. To maximize the freedom to speak and eliminate problems of gender bias, repression and domination, participants were interviewed separately by gender.

Data qualitative analysis was thematic. The transcripts of the research assistants were typed in Microsoft Excel format and processed. Afterwards the summaries of the main ideas and observations were organized according to the opinion of the majority.

The study was authorized by HSF Board and Scientific Committee, NPHD, the Secretariat of the Administrative Post of Natikiri, the Lúrio University Institutional Health Bioethics Committee and the Bioethics Committee of the University of Saskatchewan, following all Helsinki Declaration (2013) recommendations.

No changes were made in the study procedures to the initial protocol.

3. Results

300 people were surveyed (see Table I) and 11 FGD were held (see Table II).

Table I: sample of subjects interviewed.

Women of Childbearing Age	77
Male Heads of Households	78
Older Women	75
TMW and THP	46
Community Leaders	15
Maternal Child Health nurses	9
Total	300

Table II: focus groups discussions.

Women of Childbearing age	1
Male Heads of households	1
Adolescent mothers	1
Older Women	1
Community Leaders	1
TMW	2
THP	2
Maternal Newborn care nurses	2
Total	11

The answers on the questionnaire were classified using the Likert scoring system

(Always = 1, Sometimes = 2, Never = 3). The mean and standard deviation for each response was calculated and compared between groups. Consistent themes emerged across all groups and were supported by data from shared testimonies in the FGD. The consistent themes found were:

1. Limited knowledge about maternal health and family planning (FP) needs.

2. Lack of transportation to access maternal newborn health care.

3. Poor quality of maternal health care.

4. Continued need for government support and funding for maternal newborn care.

Due to the limited knowledge and access to FP women become pregnant again a very short time after delivery. Nurses (to a greater extent) and men (to a lesser extent) recognize this concern (see Table III). A maternal newborn nurse in the focus group justifies this situation:

"There are mothers who forbid contraceptives spoil the girls' daughters to continue the FP because reproductive system." they want grandchildren and say that

Table III: spacing between pregnancies.

Question: the majority of women become pregnant too soon after their last delivery? 1= always, 2 =Sometimes 3 = Never			
Group	N	\bar{x}	s.d.
Men	78	2.08	0.79
Women of childbearing age	77	1.82	0.66
Older Women	75	1.85	0.73
Community Leaders	15	1.67	0.62
Maternal Newborn nurses	9	1.33	0.50
TMW and THP	46	1.85	0.69

Another recognized problem is the high frequency of early pregnancy; a community leader in a FGD said:

"The issue of early pregnancy is worrying and those in the community do not know how to overcome this problem, but people in the community care a lot."

Women find it difficult to access health care because transport options are limited. All groups reported that women go on foot to consultations and rarely went by public transport or by car.

Unsafe travel conditions were also cited as one of the main concerns (see Table IV). A community leader participating in a FGD summarized the situation:

"They require lights on public roads and paths. This could make it easier for people to go to the hospital at night, and it would also help if a woman gives birth on the way to the hospital".

Table IV: access to care for pregnant women.

Question: Do women go to antenatal visits on foot? 1= always, 2 = sometimes 3 = never			
Group	N	\bar{x}	s.d.
Men	78	1.67	0.62
Women of Childbearing Age	77	1.41	0.61
Older Women	75	1.52	0.81
Community Leaders	15	1.40	0.57
Maternal Newborn nurses	9	1.78	0.44
TMW and THP	46	1.41	0.69

The quantitative results did not provide a clear picture of the HC experience of maternal and newborn health care. However, qualitative evidence demonstrates knowledge of specific difficulties for quality health care. A FGD participant parent said:

"When I arrived with my wife, the maternal health nurse did not greet us sympathetically and I could say she despised us, perhaps because we seemed very poor."

A THP said:

"Once there was a woman in labor and the maternal newborn nurse simply abandoned her and went to bed. When the TMW tried to help her, the maternal newborn nurse threatened to leave the woman in labor and also insulted the TMW".

One parent reinforces:

"When I arrived with my pregnant wife crying in pain, the maternal newborn nurse said to my wife laughingly: you are crying now, but what were you

thinking when you were doing these things (sex)?"

A community leader adds,

"One of the mothers said that when she was pregnant, at the time of delivery, after being admitted, the maternity nurse asked if she had any money. She asked how much, but the maternity nurse did not answer but left her alone during labor and she gave birth alone. She just called the maternity nurse to finish the rest. "

Maternal, neonatal and child health, in the opinion of the participants, is considered a priority for the Government of Mozambique. This result is evident both in the quantitative data (see Table V) and in the FGD.

Table V: how does the Government value maternal and newborn health?

Question: Maternal health is an important priority for the government 1= always 2 = sometimes 3 = never			
Group	N	\bar{x}	s.d.
Men	78	1.83	1.21
Women of Childbearing Age	77	1.62	1.00
Older Women	75	1.23	0.56
Community Leaders	15	1.27	0.46
Maternal newborn nurses	9	2.11	0.78
TMW and THP	46	1.61	1.02

But the continuing need for Government support and funding of policies for maternal and newborn care is also

recognized. A maternal newborn nurse participant in a FGD said:

"The Ministry of Health should continue to require husbands to accompany their

wives to ANC visits because FP is explained in detail in these visits."

One community leader said:

"One of the biggest problems is that TMW do not feel valued or recognized in the HC and do not receive any payment. They work but receive nothing in return and are requesting, at least, some payment."

A parent head of household said:

"Sometimes the Government provides some support to the community, like mosquito nets and meals for children or pregnant women, to combat malnutrition, however, there is a need for transparency in distribution so that community leaders and HP do not prevent supplies from reaching the community and keep them for personal gain."

The main results can be summarized in the themes that were consistently present in all FGD:

1. Long walking distances required to get to the HC or MH.
2. Unsafe travel conditions for women who walk alone.
3. Bad treatment in HC including illicit payments and bribes.
4. Long waits to be attended in the HC or MH.
5. HP poor training and knowledge.

6. Neglect and lack of family centered care.

7. Women who become pregnant very soon after their last birth or at very young age.

8. Inability of women to make informed decisions about FP.

9. Continued practices of home births and traditional methods of treatment.

10. Myths and cultural taboos about pregnancy and newborn care.

11. Limited knowledge and understanding among women and the community regarding the health needs of women and young people.

12. Need for "Maternity Waiting homes" near the hospital for pregnant women.

13. Weak Government policies and little funding to support MCH care.

FGD participants identified priorities for maternal and newborn health:

- 1) Transportation options to meet the long distances needed to reach the HC.
- 2) Lack of attendance at ANC visits and neglect of maternal care needs.
- 3) Poor and insufficient service in MH.
- 4) Lack of qualified HP in childbirth care.
- 5) The need for Government policies to support adherence to maternal care and respectful treatment for mothers.

4. Discussion

International literature review identifying factors causing delay in the search for care and treatment by pregnant and newborn confirm this study results:

18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55

1) Delay in the decision to seek care:

a) Decision to seek health care is dependent on husband.

b) Decision to use FP is dependent on husband.

c) Lack of knowledge of pregnancy warning signs.

d) Lack of basic knowledge about SRH.

e) Lack of support for household chores.

f) Lack of confidence in the health system.

g) Cultural taboos and witchcraft.

h) Poor participation of men in maternal and neonatal care.

i) High cost of health services.

2. Delay on arrival at HC:

a) Long distance to HC.

b) Lack of money for transportation payment.

c) Lack of means of transport.

d) Conditional authorization to seek health care.

e) Do not have a trusted or supportive person.

3. Delay in providing quality care

a) Hospital lacking HP.

b) Hospital lack of resources.

c) Maternal newborn nurse professional negligence.

d) Maternal newborn nurse professionals with poor training.

e) Maternal newborn nurse services not being a Government priority.

5. Conclusion

The baseline study and consultations with local community and institutional partners, indicate six intervention lines, evidence based, that might reverse the problems identified. These are:

1. Expanding FP especially with adolescents.

2. Community based transport system for pregnant women.

3. Strengthening maternal and child health services by training maternity personnel in Obstetrical Emergency care (EmOC) and neonatal resuscitation (Helping Babies Breath).

4. Providing four quality prenatal visits.

5. Providing quality cesarean deliveries.

6. Supporting the Mozambican bribery.
government's campaign against

References

- ¹ Cousens S, Blencowe H, Stanton C. e col. As estimativas nacionais, regionais e mundiais de natimortalidade em 2009, com as tendências desde 1995. Uma análise sistemática. Lancet 2011, 377 (9774): 1319-1330.
- ² WHO. International statistical classification of diseases and related health problems, tenth revision. Vol. 1: Tabular list. Vol. 2: Instruction manual. World Health Organization. Geneva. 2010.
- ³ Say L. e col. Causas globais de Morte Materna: Uma análise sistemática. OMS Lancet. 2014.
- ⁴ Ministério da Planificação e Desenvolvimento. Relatório Sobre os Objectivos de Desenvolvimento do Milénio de 2010. Governo de Moçambique, Instituto Nacional de Estatística e Estatísticas dos Sectores. República de Moçambique. Maputo. 2010. www.mpd.gov.mz.
- ⁵ Departamento de Saúde da Comunidade, Secção de Saúde Infantil. Política Nacional de Saúde Neonatal e Infantil em Moçambique. Direcção Nacional de Saúde. Ministério da Saúde. República de Moçambique. Maputo, 2006.
- ⁶ Instituto Nacional de Estatística. Moçambique, Inquérito Demográfico e de Saúde 2011. Relatório Preliminar. Ministério da Saúde. Maputo, 2012.
- ⁷ MISAU. Estratégia para o Fortalecimento das Intervenções da Parteira Tradicional. Ministério da Saúde. República de Moçambique. Maputo. 2009. www.misau.gov.mz.
- ⁸ A Biza, I Jille-Traas, M Colomar e col. Challenges and opportunities for implementing evidence-based antenatal care in Mozambique: a qualitative study. BMC Pregnancy and Childbirth (2015) 15:200.
- ⁹ P. Garrido, A. Libombo, M. Saide. Roteiro para acelerar a redução da mortalidade materna e neonatal em Moçambique. Ministério da Saúde, República de Moçambique. Maputo. 2008.
- ¹⁰ S. Rokia, S. Giani. Valorisation du rôle des accoucheuses traditionnelles dans la prise en charge des urgences obstétricales au Mali. Ethnopharmacologia, n°43, DOSSIER SPÉCIAL : Médecine traditionnelle en Afrique. Juillet 2009.
- ¹¹ UNICEF. Situação das Crianças em Moçambique 2014. Maputo. 2014. ISBN: 978-92-806-4769-3. UNICEF. http://sitan.unicef.org.mz/files/UNICEF_FULL_Situacao-das-Crianças-em-Mocambique_Portugues.pdf.

- ¹² Lawn E, Cousens S, Zupan K. Lancet Neonatal Survival Steering Team. 4 million neonatal deaths: When? Where? Why? *The Lancet* 2005;365(9462):891-900.
- ¹³ WHO. Health and the Millennium Development Goals. World Health Organization. Geneva. 2005.
- ¹⁴ Ministério da Saúde. Caderneta de Saúde da Mulher. World Health Organization. Maputo. 2011.
- ¹⁵ WHO. Increasing access for child and maternal health care services: the Mozambique experience. WHO/AFRO Library Cataloguing – in – Publication. WHO Regional Office for Africa. 2013.
- ¹⁶ L. Mavota, G. Snetro, C. Regina. Termos de referência para o estabelecimento e funcionamento dos Comitês de Gestão das Unidades Sanitárias. Ministério da Saúde. Direcção Nacional de Saúde. Maputo. 2011.
- ¹⁷ P. Garrido, A. Libombo, M. Saíde. Iniciativa Maternidades modelo: padrões para medição do desempenho dos serviços de saúde materna e neonatal. Ministério da Saúde. República de Moçambique. Maputo. 2009.
- ¹⁸ E. Abalos, E. Chamillard, V. Diaz e col. Antenatal care for healthy pregnant women: a mapping of interventions from existing guidelines to inform the development of new WHO guidance on antenatal care, 2016. *BJOG*. 123(4):519-28.
- ¹⁹ S. Ahmed, M. Norton, E. Williams e col. Operations research to add postpartum family planning to maternal and neonatal health to improve birth spacing in Sylhet District, Bangladesh, 2013. *Global Health Science Practice*. 141(2):262-76.
- ²⁰ B. Hamdela, A. Mariam, T. Tilahun. Unwanted Pregnancy and Associated Factors among Pregnant Married Women in Hosanna Town, Southern Ethiopia, 2012. *PLoS One*. 7(6).
- ²¹ L. Benova, O. Cumming, B. Gordon e col. Where There Is No Toilet: Water and Sanitation Environments of Domestic and Facility Births in Tanzania, 2014. *PLOS One*. 9(9).
- ²² A. Canavan. Review of global literature on maternal health interventions and outcomes related to provision of skilled birth attendance. Royal Tropical Institute. Amsterdam. 2009.
- ²³ L. Chola, S. McGee, A. Tugendhaft e col. Scaling Up Family Planning to Reduce Maternal and Child Mortality: The Potential Costs and Benefits of Modern Contraceptive Use in South Africa, 2015. *PLoS One*. 10(6).

- ²⁴ E. Duysburgh, B. Kerstens, S. Kouanda e col. Opportunities to improve postpartum care for mothers and infants: design of context-specific packages of postpartum interventions in rural districts in four sub-Saharan African countries, 2015. *BMC Pregnancy Childbirth*. 3(15):131.
- ²⁵ S. Eliason, F. Baiden, G. Quansah-Asare e col. Factors influencing the intention of women in rural Ghana to adopt postpartum family planning, 2013. *Reproductive Health*.10:34.
- ²⁶ M. Ellsberg. Intimate partner violence and women's physical and mental health in the WHO multi-country study on women's health and domestic violence: an observational study, 2008. *The Lancet* 371.9619: 1165-1172.
- ²⁷ A. Exavery, A. Kanté, N. Mustafa e col. Access to institutional delivery care and reasons for home delivery in three districts of Tanzania, 2014. *International Journal for Equity in Health*.13:48.
- ²⁸ E. Falnes, K. Moland, T. Tylleskär e col. It is her responsibility: partner involvement in prevention of mother to child transmission of HIV programs, northern Tanzania, 2011. *Journal of the International AIDS Society*.14:21.
- ²⁹ C. García-Moreno. WHO multi-country study on women's health and domestic violence against women. World Health Organization. Geneva. 2005.
- ³⁰ N. Gerein, A. Green, S. Pearson. The implications of shortages of health professionals for maternal health in sub-Saharan Africa, 2006. *Reproductive Health Matters*. 14(27):40-50.
- ³¹ K. Gross, I. Mayumana, B. Obrist. My wife, you are supposed to have a rest now: an analysis of norms influencing men's role in prenatal care in south-eastern Tanzania, 2013. *Anthropology & Medicine*. 20:1.
- ³² S. Gupta, G. Yamada, R. Mpembeni e col. Factors Associated with Four or More Antenatal Care Visits and Its Decline among Pregnant Women in Tanzania between 1999 and 2010. 2014. *PLoS One*. 9(7).
- ³³ A. Kearns, J. Caglia, P. Hoope-Bender e col. Antenatal and postnatal care: a review of innovative models for improving availability, accessibility, acceptability and quality of services in low-resource settings. 2016. *BJOG*. 123(4):540-8.
- ³⁴ A. Kes, S. Ogwang, R. Pande e col. The economic burden of maternal mortality on households: evidence from three sub-counties in rural western Kenya. 2015. *Reproductive Health*. 6(12):1.
- ³⁵ S. Kujawski, G. Mbaruku, L. Freedman e col. Association Between Disrespect and Abuse During Childbirth and Women's Confidence in Health Facilities in Tanzania. 2015. *Maternal Child Health Journal*. 19(10):2243-50.

- ³⁶ A. Luz, M. Osis, M. Ribeiro e col. Impact of a nationwide study for surveillance of maternal near-miss on the quality of care provided by participating centers: a quantitative and qualitative approach. 2014. BMC Pregnancy Childbirth.1(14):122.
- ³⁷ M. Magoma, J. Requejo, M. Meriardi e col. How much time is available for antenatal care consultations? Assessment of the quality of care in rural Tanzania. 2014. BMC Pregnancy and Childbirth. 11:64.
- ³⁸ A. Mazzoni, F. Althabe, L. Gutierrez e col. Women's preferences and mode of delivery in public and private hospitals: a prospective cohort study. 2016. BMC Pregnancy and Childbirth.16:34.
- ³⁹ M. Meskele, W. Mekonnen. Factors affecting women's intention to use long acting and permanent contraceptive methods in Wolaita Zone, Southern Ethiopia: A cross-sectional study. 2014. BMC Women's Health.14:109.
- ⁴⁰ G. Molina, T. Weiser, S. Lipsitz e col. Relationship Between Cesarean Delivery Rate and Maternal and Neonatal Mortality. 2015. Jama 1:314(21).
- ⁴¹ L. Mselle, T. Kohi, A. Mvungi e col. Waiting for attention and care: birthing accounts of women in rural Tanzania who developed obstetric fistula as an outcome of labour. 2011. BMC Pregnancy and Childbirth. 11:75.
- ⁴² L. Mselle, K. Moland, A. Mvungi e col. Why give birth in health facility? Users' and providers' accounts of poor quality of birth care in Tanzania. 2013. BMC Health Services Research. 13:174.
- ⁴³ R. Muela, S. Hausmann-Muela. The straw that breaks the camel's back. 2011. Medical Anthropology Quarterly. 25(1):103–21
- ⁴⁴ K. Rajendra, C. Binns, A. Lee. Determinants of facility delivery after implementation of safer mother programme in Nepal: a prospective cohort study. 2013. BMC Pregnancy and Childbirth. 13:193.
- ⁴⁵ F. Richard, S. Zongo, F. Ouattara. Fear, guilt, and debt: an exploration of women's experience and perception of cesarean birth in Burkina Faso, West Africa. 2014. International Journal of Women's Health. 6: 469–478.
- ⁴⁶ L. Sibley, M. Tesfaye, S. Desta e col. Improving Maternal and Newborn Health Care Delivery in Rural Amhara and Oromiya Regions of Ethiopia Through the Maternal and Newborn Health in Ethiopia Partnership. 2014. Journal of Midwifery & Women's Health. 59: S6–S20.
- ⁴⁷ A. Srivastava, B. Avan, P. Rajbangshi e col. Determinants of women's satisfaction with maternal health care: a review of literature from developing countries. 2015. BMC Pregnancy and Childbirth. 15:97.

-
- ⁴⁸ Y. Tebekaw, B. Aemro, C. Teller. Prevalence and determinants of unintended childbirth in Ethiopia. 2014. BMC Pregnancy and Childbirth. 14:326.
- ⁴⁹ F. Tesfahun, W. Worku, F. Mazengiya e col.. Knowledge, perception and utilization of postnatal care of mothers in Gondar Zuria District, Ethiopia: a cross-sectional study. 2014. Maternal Child Health Journal. 18(10):2341-51.
- ⁵⁰ P. Wangwe, M. Nyasinda, D. Charles. Effectiveness of counseling at primary health facilities: Level of knowledge of antenatal attendee and their attitude on Prevention of Mother to Child Transmission of HIV in primary health facilities in Dar es salaam, Tanzania. 2014. African Health Sciences.14(1): 150-156.
- ⁵¹ A. Worku, G. Tessema, A. Zeleke. Trends of Modern Contraceptive Use among Young Married Women Based on the 2000, 2005, and 2011 Ethiopian Demographic and Health Surveys: A Multivariate Decomposition Analysis. 2015. PLoS One. 10(1).
- ⁵² E. Yesuf, R. Calderon-Margalit. Disparities in the use of antenatal care service in Ethiopia over a period of fifteen years. 2013. BMC Pregnancy and Childbirth.13:131.
- ⁵³ K. Finlayson, S. Downe. Why Do Women Not Use Antenatal Services in Low- and Middle-Income Countries? A Meta-Synthesis of Qualitative Studies. PLOS Medicine, January 2013, Volume 10, Issue 1. e1001373. www.plosmedicine.org.
- ⁵⁴ V. Agadjanian, J. Yao, S. Hayford. Spatial, Social, and Institutional Determinants of Child Delivery Place in Rural Mozambique. Center for Population Dynamics. Arizona State University. 2012.
- ⁵⁵ J. Driessen, Z. Dodson, V. Agadjanian, The Effects of Distance and Quality on Uptake of Sexual, Reproductive, and Other Health Services in Rural Mozambique. PFRH Seminar. December 10, 2014.
- ⁵⁶ P. Pires, R. Siemens, D. Joao e col. Women's perceptions about ante – natal care access, Marrere hospital, Nampula, Mozambique, 2014. International Journal of Research, Volume 03 Issue 14, October 2016. <https://edupediapublications.org/journals>.